

be helped by the introduction of high-throughput multiplex genotyping, which will enable simultaneous sequencing and measuring copy numbers of hundreds of genes from only nanograms of cancer cell DNA.

Although these advances will probably reduce the proportion of patients who need chemotherapy, it will remain the main treatment option for advanced NSCLC. The TAILOR study will contribute to more rational use of chemotherapy and EGFR tyrosine kinase inhibitors in the treatment of NSCLC, based on its molecular profile.

**Jacek Jassem, Rafał Dziadziuszko*

Department of Oncology and Radiotherapy, Medical University of Gdańsk, 80 211 Gdańsk, Poland
jjassem@gumed.edu.pl

We declare that we have no conflicts of interest.

- 1 Shepherd FA, Rodrigues Pereira J, Ciuleanu T, et al. Erlotinib in previously treated non-small-cell lung cancer. *N Engl J Med* 2005; **353**: 123–32.
- 2 Lynch TJ, Bell DW, Sordella R, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2004; **350**: 2129–39.

- 3 Paez JG, Jänne PA, Lee JC, et al. EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 2004; **304**: 1497–500.
- 4 Rosell R, Carcereny E, Gervais R, et al. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 2012; **13**: 239–46.
- 5 Mitsudomi T, Morita S, Yatabe Y, et al. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. *Lancet Oncol* 2010; **11**: 121–28.
- 6 Zhou C, Wu YL, Chen G, et al. Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre, open-label, randomised, phase 3 study. *Lancet Oncol* 2011; **12**: 735–42.
- 7 Gerassino MC, Martelli O, Broggin M, et al. A randomised trial of erlotinib versus docetaxel as second-line treatment of patients with advanced non-small cell lung cancer and wild-type epidermal growth factor receptor (TAILOR). *Lancet Oncol* 2013; published online July 22. [http://dx.doi.org/10.1016/S1470-2045\(13\)70310-3](http://dx.doi.org/10.1016/S1470-2045(13)70310-3).
- 8 Maheswaran S, Sequist LV, Nagrath S, et al. Detection of mutations in EGFR in circulating lung-cancer cells. *N Engl J Med* 2008; **359**: 366–77.
- 9 Bai H, Mao L, Wang HS, et al. Epidermal growth factor receptor mutations in plasma DNA samples predict tumor response in Chinese patients with stages IIIB to IV non-small-cell lung cancer. *J Clin Oncol* 2009; **27**: 2653–59.
- 10 Li T, Kung HJ, Mack PC, Gandara DR. Genotyping and genomic profiling of non-small-cell lung cancer: implications for current and future therapies. *J Clin Oncol* 2013; **31**: 1039–49.

Combination endocrine treatments unproven in breast cancer




Ovarian ablation, which was introduced more than 100 years ago, was the first endocrine treatment for advanced breast cancer, followed by adrenalectomy and hypophysectomy. These ablative therapies have since been replaced by antioestrogen treatments, luteinising-hormone-releasing hormone agonists, and aromatase inhibitors.¹ Other endocrine treatments with different mechanisms of action have also become available for breast cancer: oestrogens, progestins, androgens, antiandrogens, and selective oestrogen-receptor downregulators.

Although the superiority of combinations of chemotherapeutic agents with different mechanisms of action to single agents has been established in the treatment of early and advanced breast cancer,² that of combinations of endocrine treatments has not been shown.¹ Several combinations of hormonal agents have been assessed in patients with advanced breast cancer, with no consistent improvement in either time to progression of disease or survival.³ Additionally, the combination of an antioestrogen treatment (tamoxifen) and ovarian suppression with a luteinising-hormone-releasing hormone agonist does not lead to improvements in disease-free or

overall survival when compared with antioestrogen treatment alone in premenopausal women with early breast cancer.⁴ A large, prospective, double-blind trial⁵ comparing tamoxifen with an aromatase inhibitor (anastrozole) and tamoxifen or anastrozole alone as adjuvant treatment for breast cancer showed that the combination did not improve either disease-free or overall survival. Indeed, the group who received the combination was discontinued after initial analysis of the data.⁵

Because preclinical data⁶ suggested that the combination of an aromatase inhibitor and fulvestrant—a member of the newest class of hormonal agents, selective oestrogen receptor downregulators—was superior to either agent alone against breast tumours in mice, three prospective studies^{7–9} have assessed this approach in postmenopausal women with advanced breast cancer. In *The Lancet Oncology*, Stephen Johnston and colleagues report results of a phase 3, European, multicentre trial.⁷ Postmenopausal women with hormone-receptor-positive breast cancer who had relapsed or progressed while receiving a non-steroidal aromatase inhibitor were randomly assigned to receive fulvestrant plus anastrozole, fulvestrant plus



NCI Science Photo Library

Published Online
 July 29, 2013
[http://dx.doi.org/10.1016/S1470-2045\(13\)70364-4](http://dx.doi.org/10.1016/S1470-2045(13)70364-4)

See [Articles](#) page 989

Copyright © Buzzdar. Open Access article distributed under the terms of CC BY-NC-SA

anastrozole-matched placebo, or the steroidal aromatase inhibitor exemestane. No improvement in progression-free survival was recorded in the group who received fulvestrant plus anastrozole (median 4.4 months, 95% CI 3.4–5.4) compared with fulvestrant plus placebo (4.8 months, 3.6–5.5; hazard ratio [HR] 1.00, 95% CI 0.83–1.21; log-rank $p=0.98$), or in the group given fulvestrant plus placebo compared with exemestane (3.4 months, 3.0–4.6; HR 0.95, 0.79–1.14; log-rank $p=0.56$).

Another phase 3 study⁸ comparing the combination of anastrozole and fulvestrant as first-line treatment in postmenopausal patients with hormone-receptor-positive advanced breast cancer also showed no advantages in terms of clinical efficacy for the combination compared with anastrozole alone. By contrast, a phase 3 trial from North America⁹ compared the combination of anastrozole and fulvestrant with anastrozole alone as first-line treatment in postmenopausal women with hormone-receptor-positive advanced breast cancer, and showed increased control of disease and survival with the combination of fulvestrant and anastrozole. However, the apparent improvements could have been due to imbalances in prognostic characteristics between the two study groups.

Therefore, there are still no consistent data to support the notion that combination endocrine therapy is superior to treatment with one agent in early or advanced breast cancer. However, as understanding of the mechanisms of resistance to endocrine treatment has improved, targeting of some pathways has resulted in new approaches that offer hope for disease control. For example, postmenopausal patients with hormone-receptor-positive, HER2-positive advanced breast cancer given endocrine and anti-HER2 treatments have had longer control of disease than have those given endocrine treatment alone.¹ Indeed, a combination of an aromatase inhibitor with an anti-HER2 treatment has been approved by the US Food and Drug Administration for the management of postmenopausal patients with hormone-receptor-positive, HER2-positive advanced breast cancer.¹ Another effective approach has been the combination of an mTOR inhibitor (everolimus) with exemestane.¹⁰ The combination of exemestane with a histone deacetylase inhibitor (entinostat) has also had encouraging results.¹¹

In conclusion, a combination of endocrine agents with different mechanisms of action will probably not result in a meaningful improvement in outcomes for patients with breast cancer. Sequential use of endocrine agents remains the standard of care in patients with advanced breast cancer. However, understanding of the mechanisms of resistance to hormonal agents continues to advance, and combinations of endocrine treatment with targeted agents that block resistance pathways should improve the outlook for patients with breast cancer that is resistant to endocrine treatment.

Aman U Buzdar

Department of Breast Medical Oncology, University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA
abuzdar@mdanderson.org

I declare that I have no conflicts of interest.

- 1 Barrios C, Forbes JF, Jonat W, et al. The sequential use of endocrine treatment for advanced breast cancer: where are we? *Ann Oncol* 2012; **23**: 1378–86.
- 2 Early Breast Cancer Trialists' Collaborative Group. Effects of adjuvant tamoxifen and of cytotoxic therapy on mortality in early breast cancer: an overview of 61 randomized trials among 28 896 women. *N Engl J Med* 1988; **319**: 1681–92.
- 3 Michaud LB, Jones KL, Buzdar AU. Combination endocrine therapy in the management of breast cancer. *Oncologist* 2001; **6**: 538–46.
- 4 LHRH-agonists in Early Breast Cancer Overview group. Use of luteinising-hormone-releasing hormone agonists as adjuvant treatment in premenopausal patients with hormone-receptor-positive breast cancer: a meta-analysis of individual patient data from randomised adjuvant trials. *Lancet* 2007; **369**: 1711–23.
- 5 The ATAC (Arimidex, Tamoxifen Alone or in Combination) Trialists' Group. Anastrozole alone or in combination with tamoxifen versus tamoxifen alone for adjuvant treatment of postmenopausal women with early breast cancer: first results of the ATAC randomised trial. *Lancet* 2002; **359**: 2131–39.
- 6 Jelovac D, Macedo L, Golubeva OG, Handratta V, Brodie AM. Additive antitumor effect of aromatase inhibitor letrozole and antiestrogen fulvestrant in a postmenopausal breast cancer model. *Cancer Res* 2005; **65**: 5439–44.
- 7 Johnston SRD, Kilburn LS, Ellis P, et al. Fulvestrant plus anastrozole or placebo versus exemestane alone after progression on non-steroidal aromatase inhibitors in postmenopausal patients with hormone-receptor-positive locally advanced or metastatic breast cancer (SoFEA): a composite, multicentre, phase 3 randomised trial. *Lancet Oncol* 2013; published online July 29. [http://dx.doi.org/10.1016/S1470-2045\(13\)70322-X](http://dx.doi.org/10.1016/S1470-2045(13)70322-X).
- 8 Bergh J, Jönsson PE, Lidbrink EK, et al. FACT: an open-label randomized phase III study of fulvestrant and anastrozole in combination compared with anastrozole alone as first-line therapy for patients with receptor-positive postmenopausal breast cancer. *J Clin Oncol* 2012; **30**: 1919–25.
- 9 Mehta RS, Barlow WE, Albain KS, et al. Combination anastrozole and fulvestrant in metastatic breast cancer. *N Engl J Med* 2012; **367**: 435–44.
- 10 Baselga J, Campone M, Piccart M, et al. Everolimus in postmenopausal hormone-receptor-positive advanced breast cancer. *N Engl J Med* 2012; **366**: 520–29.
- 11 Yardley DA, Ismail-Khan RR, Melichar B, et al. Randomized phase II, double-blind, placebo-controlled study of exemestane with or without entinostat in postmenopausal women with locally recurrent or metastatic estrogen receptor-positive breast cancer progressing on treatment with a nonsteroidal aromatase inhibitor. *J Clin Oncol* 2013; **31**: 2128–35.